

Interacting Quintessence, the Coincidence Problem and Cosmic Acceleration

Greg Huey¹ and Benjamin D. Wandelt^{1,2}

¹*Department of Physics, University of Illinois, Urbana, IL 61801*

²*Department of Astronomy, UIUC, 1002 W Green Street, Urbana, IL 61801*

Faced by recent evidence for a flat universe dominated by dark energy, cosmologists grapple with deep cosmic enigmas such as the cosmological constant problem, extreme fine-tuning and the cosmic coincidence problem. The extent to which we observe the dimming of distant supernovae suggests that the cosmic acceleration is as least as severe as in cosmological constant models. Extrapolating this to our cosmic future implies terrifying visions of either a cold and empty universe or an explosive demise in a “Big Rip.” We construct a class of dynamical scalar field models of dark energy and dark matter. Within this class we can explain why supernovae imply a cosmic equation of state $w \lesssim -1$, address fine tuning issues, protect the universe from premature acceleration and predict a constant fraction of dark energy to dark matter in the future (thus solving the coincidence problem), satisfy the dominant energy condition, and ensure that gravitationally bound objects remain so forever (avoid a Big Rip). This is achieved with a string theory inspired Lagrangian containing standard kinetic terms, exponential potentials and couplings, and parameters of order unity.

There is mounting evidence for the presence of an enigmatic dark energy component of the Universe. The nature of this component is one of the most profound mysteries of physics (see [1] for a recent review). Dark energy was first implied by a measurement of the luminosity distance-redshift relation, $d_L(z)$, through observations of supernova Ia (SNIa) at cosmological distances [2, 3]. Recent SNIa data, galaxy redshift catalogues and analysis of the cosmic microwave background anisotropy favor a cosmic dark energy equation of state (w_Q) close to or even below -1 [4, 5, 6, 7].

Models in which this missing energy is a cosmological constant Λ fit this data well. However, these models have two serious drawbacks: the fine-tuning problem (Λ must be fine-tuned extremely precisely in the early universe relative to the energy scale at that time); and the coincidence problem (Λ must be set to an extremely small value in the early universe relative to the energy scale at that time such that matter domination and structure formation could occur). Cosmological constant dark matter (Λ CDM) models are therefore not an explanation but at best an effective description of the dark energy.

Simple models of dark energy as dynamical, non-interacting scalar fields, “quintessence” [8], do not require fine-tuning of energy densities or field values at early times, because the field follows an attractor solution as it rolls down a potential [9, 10]. However, the potential must still yield a ratio of matter to quintessence energy density of $\rho_{M0}/\rho_{Q0} \sim 1/2$ today, while allowing a matter dominated phase so that structure can form. Such models therefore do not address the coincidence problem. Furthermore, there is difficulty achieving a current quintessence equation of state $\lesssim -0.8$ while retaining significant matter density [10], except with quintessence potentials designed specifically for that purpose [11].

The possibility that $w_Q < -1$, i.e. that the sum of pressure and energy density of the dark energy is less than zero, has led to the idea of “phantom energy” [12]. Explicit scalar field models of phantom energy produce super-exponential expansion of the universe by introduc-

ing a negative kinetic term in the Lagrangian. The scale factor reaches infinity in finite time and the universe ends in an explosive “Big Rip.” Setting aside theoretical difficulties of such models, e.g. the fact that they can lead to unstable solutions by virtue of violating the Dominant Energy Condition [13], the cosmic coincidence is not explained. In fact we would argue that the fine tuning problem is exacerbated compared to cosmological constant models. Instead one appeals to an anthropic argument: if conscious life appears at all it must do so within the brief era of cosmic structure between the onset of matter domination and the beginning of cosmic acceleration since the universe is destroyed soon after [14].

We look upon these difficulties with simple models as opportunities—guideposts in the vastness of theory-space to direct our attention. A natural expectation in the context of string-inspired cosmologies are interacting fields in the dark sector (see [15] and references therein). Whilst the simplest models are tentatively ruled out by comparison to cosmic microwave background observations [16], previous studies did not consider the possibility of more than one dark matter component. In this letter, we generalize the class of interacting dark matter and dark energy models suggested in [15, 17] and demonstrate by means of an explicit example that this resolves the above difficulties.

We call this class interacting quintessence cold dark matter (iQCDM) models. Our generalization consists in imagining additional field content of the dark sector. We propose that there is a quintessence-like rolling self-coupled scalar field ϕ as well as other fields that can act as dark matter. These dark matter components will generically be coupled to ϕ , but the details of the couplings would depend on the details of the underlying fundamental theory. In general one of the components will have a more ϕ dependent coupling than the others during the brief period of cosmological history between matter radiation equality and today. We call this dark matter component the interacting cold dark matter (iCDM) and group all the other components under the term non-

interacting dark matter (nCDM).

Let the quintessence field ϕ be contained in an effective potential that is the sum of two parts: a self-interaction term $V_Q(\phi) = f(\phi)$, and interaction with the iCDM, $V_M(\phi) = g(\phi)e^{-3N}$. The exponential decay (N is the logarithm of the scale factor) is a consequence of the dilution of the dark matter with expansion, an important fact we will return to below. At early times the form of the potential should prevent quintessence domination through an attractor solution (*tracking*), while at late times ϕ will be at the minimum of the potential, which should yield a supernovae-measured $w_Q \lesssim -0.9$. This yields the following conditions:

$$\frac{f''f}{(f')^2}, \frac{g''g}{(g')^2} \geq 1; \quad \frac{-f'}{\kappa f}, \frac{-g'}{\kappa g} \gtrsim 5; \quad \frac{d\phi}{dN}\kappa = \frac{3\kappa}{\frac{f''}{f'} + \frac{g''}{g'}} \lesssim \frac{1}{2},$$

where N is expansion e-folds and $\kappa \equiv \sqrt{8\pi G}$. These conditions arise from: a requirement that tracking solutions exist [10], that the tracking solutions keep ϕ subdominant [9] and that once at the potential minimum ϕ will grow to dominate with an apparent $w_Q \lesssim -0.9$. This apparent w_Q is what supernovae measurements would favor given a measurement of Ω_{M0} based on the assumption that matter did not interact with the quintessence. Steeply curved functions are favored by the second and third conditions. An additional criterion is that a significant period ($\gtrsim 7$ e-folds) of matter domination precede the onset of quintessence domination. For a specific implementation one would like to find simple forms for f and g which satisfy these conditions and arise naturally in the low-energy limit of string theory. While this may seem to be a tall order we will now go on to show that we can simultaneously satisfy all these constraints using exponential potentials and couplings.

As an illustrative example, consider the following specific model:

$$\mathcal{L} \supset \frac{\dot{\phi}^2}{2} + \frac{\dot{\chi}^2}{2} - m_Q^4 e^{-\alpha\kappa\phi - \gamma} \left(e^{\beta\kappa(\phi - \phi_c)} \right)^2 \chi^2 + \mathcal{L}_{nCDM} + \mathcal{L}_B,$$

where the quintessence field ϕ interacts with a scalar iCDM field χ with coupling constant γ , $\kappa \equiv \sqrt{8\pi G}$, $\mathcal{L}_{nCDM}$ contains nCDM, and $\mathcal{L}_B$ contains baryons. The constants α , β , γ , m_Q , ϕ_c would take values from an underlying physical theory. The value of ϕ minus some offset ϕ_c sets the mass of χ . The magnitude of the mass scale m_Q is not actually important—it can be changed to current quintessence scales by a shift of ϕ and ϕ_c .

One can “integrate out” χ if it oscillates on a much faster timescale than the change of ϕ . This calculation [22] results in an effective potential V , including the exponentially decaying interaction term, and the following equation of motion for ϕ :

$$\begin{aligned} \frac{d^2\phi}{dt^2} + 3H\frac{d\phi}{dt} + V_{,\phi} &= 0 & V &\equiv V_Q + V_M \\ V_Q &\equiv m_Q^4 e^{-\alpha\kappa(\phi)} & & \text{(quintessence)} \\ V_M &\equiv \rho_{M0} \frac{e^{\beta\kappa(\phi - \phi_c) + r}}{e^{\beta\kappa(\phi_0 - \phi_c) + r}} e^{3(N_0 - N)} & & \text{(all matter)} \\ \rho &= \rho_R + \frac{1}{2} \left(\frac{d\phi}{dt} \right)^2 + V_Q + V_M & & \text{(total energy density)}. \end{aligned}$$

($N_0 - N$) is the number of expansion e-folds before the present. The effective Lagrangian leading to these equations resembles that of exponential VAMP models [18], except for the crucial difference of including additional species of dark matter (nCDM). We effectively model the impact of the baryonic matter and nCDM on the evolution of ϕ through the term $r/e^{\beta\kappa(\phi_0 - \phi_c)}$, which sets the ratio of this non-interacting matter to iCDM today. From the point of view of ϕ , non-interacting species of matter have the same effect as the offset ϕ_c in the interaction potential. Thus, without loss of generality, one can absorb the magnitude of r into a shift of ϕ_c . For simplicity we shall set $r = 1$.

A schematic representation of the evolution of ϕ in its effective potential V is shown in Fig. 1. The potential V is the sum of a positive and negative steep exponential. Between them is a non-stationary minimum. We show the evolution of the fractional densities of radiation, matter, and dark energy for a numerical example in Fig. 2. At early times, ϕ will be on one side of the potential, high above the minimum. To the left of the minimum, ϕ will track down the potential, always remaining a fixed, small portion of the total energy density, $\rho_\phi/\rho_{tot} = 3(w_{BG} + 1)/\alpha^2$ [9], until it reaches the minimum. The background equation of state w_{BG} is $\frac{1}{3}$ during radiation domination, and 0 after matter dominates. Primordial light-element abundance measurements constrain the total energy density at the epoch of Big Bang Nucleosynthesis, and thus we must require $\alpha \gtrsim 5$ [19]. On the right-hand side of the minimum, ϕ will similarly track down this decaying potential to the minimum, keeping a fixed, small portion of the total energy density: $\rho_\phi/\rho_{tot} = 3w_{BG}/2\beta^2$ [20] until matter domination or until it reaches the minimum. At early times this tracking keeps quintessence largely independent of initial conditions and protects the universe from premature acceleration. Note that iCDM domination is also not possible while tracking down the right side of the potential because there is no tracking solution to the field equations for $w_{BG} \leq 0$.

Once ϕ reaches the minimum of the effective potential, ϕ_{min} , it evolves with it. ϕ_{min} increases with expansion because V_M is decaying—that is, the interacting dark matter is diluted by expansion. ϕ can only roll down V_Q as fast as ϕ_{min} evolves, and for steep potentials it acquires a potential dominated state with the following properties:

$$\begin{aligned} \frac{V_Q}{V_M} \Big|_{min} &= \frac{\beta}{\alpha} \frac{e^{\beta\kappa(\phi - \phi_c)}}{e^{\beta\kappa(\phi_0 - \phi_c) + 1}}; & \frac{V_Q}{\rho_{iCDM}} &= \frac{\beta}{\alpha}; \\ \kappa\phi_{min} &= \frac{3}{\alpha + \beta} N + const; & \frac{d\kappa\phi_{min}}{dN} &= \frac{3}{\alpha + \beta}; \text{ and} \\ w_{tot} + 1 &= \frac{\rho_R}{\rho} \left(\frac{4}{3} \right) + \frac{\rho_Q}{\rho} \left(\frac{\rho}{\rho_Q} \frac{3}{(\alpha + \beta)^2} \right) + \frac{\rho_M}{\rho} (1), \end{aligned}$$

where $\rho_Q \equiv \rho - \rho_R - \rho_M$ is defined as the quintessence energy density, and $\rho_M \equiv V_M$ the total matter energy density.

Therefore ϕ is in the minimum today, evolving to dominance with a current equation of state $w_{Q0} = -1 +$




Figure 1: The dashed line represents the ϕ self-interaction potential, the dot-dashed line the effective potential from the interaction with iCDM (which decays with expansion), and the solid the sum. At early times ϕ tracks down either side (either A_1 or A_2), coming to and then slowly rolling with the minimum (B).

$3/\Omega_{Q0}(\alpha + \beta)^2$. In the future, the Universe is asymptotically approaching a state where baryons and nCDM have been diluted away leaving only quintessence and interacting dark matter in the ratio $\rho_M = \rho_{iCDM} = \frac{\alpha}{\beta}\rho_Q$. Note that iCDM becoming significant near the onset of quintessence domination is not an arranged coincidence, but rather an unavoidable feature of this model due to the interactions. When ϕ is at the potential minimum, as it is during quintessence domination, the interaction guarantees that the quintessence and iCDM densities are comparable.

One might be tempted to omit the degrees of freedom which we label nCDM [18]. However, without dilution by nCDM, the energy pumped into the iCDM causes the matter to dominate over many fewer e-folds and less time and would drastically alter the matter power spectrum (since matter-radiation equality would be too recent). Furthermore, the ratio β/α would be set by the present ratio of energy in CDM to baryons ($\sim 5 : 1$) to approximately $6\Omega_{Q0}/5\Omega_{M0}$. In order to have critical density at the present temperature of the cosmic microwave background, α and β would have to be tuned to values that imply a w_Q today inconsistent with observations. Our model just requires an inequality between them to be satisfied. The amount of nCDM needed to dilute the energy pumped into iCDM depends on the value of ϕ_c , but the amount needed to allow β/α to vary is set by $\rho_{nCDM} \sim (1 - 6\alpha\Omega_{Q0}/5\beta\Omega_{M0})\rho_{CDM}$.

For our model to address the coincidence problem, achieving a constant ratio of dark energy to dark matter densities in the future is not sufficient. It must be “natural” for quintessence to dominate only after matter-radiation equality, though V_Q may be due to physics at much higher scale. This yields a mild lower limit on $\kappa\phi_c$: $\kappa\phi_c > \frac{1}{\alpha} \ln(m_Q^4/\rho_{eq})$. For $m_Q \sim TeV$, $\kappa\phi_c > 108/\alpha$. Large α reduces the size of the required ϕ_c and is also desirable for a large negative late-time quintessence equa-




Figure 2: The fraction of the energy density in radiation, matter and quintessence as a function of the natural log of the scale factor N . The thin dashed line is the tracking attractor, while the thin dotted line represents the minimum of the effective potential. Note ϕ follows the former until it crosses the latter. For this numerical example $\alpha = 5, \beta = 15, \Omega_{Q0} = 0.6$. Note that in the future, the ratio of the dark matter and dark energy densities approaches the constant α/β .

tion of state. Thus we see that the same feature of this model—large α, β with $\beta > 2\alpha$ —simultaneously avoids early cosmic acceleration both while ϕ is tracking and in the minimum of V , as well as yielding better agreement with $d_L(z)$ inferred from SNIa data, as we shall show now.

For connecting to observations it is useful to be able to talk about the quintessence and the iCDM component separately. Due to the coupling, it is somewhat arbitrary how to split the ϕ energy density into the quintessence part and the interacting dark matter part. The choice will lead to correspondingly different pressures in these components. Of course, for predictions of actual observables the choice does not have any effect. Thus, for comparison with non-interacting quintessence-CDM models, we use the split implied by the above expression for w_{tot} . With this convention, the equation of state of quintessence today can be arbitrarily close to that of a cosmological constant. Alternatively, if instead of pressure one uses the decay rates as a guide for how to split quintessence and iCDM and assign equations of state, then both quintessence and iCDM have $w = -\beta/(\alpha + \beta)$, while the non-interacting matter has $w = 0$. Since SNIa data probes $d_L(z)$, an integral of the scale factor, it ‘sees’ the iCDM as if it were part of the quintessence, both decaying as if $w = -\beta/(\alpha + \beta)$. Measurements of the matter density based on gravitational dynamics ‘see’ ρ_{iCDM} as matter. Thus SNIa and large-scale structure measurements would see a different Ω_{M0} in this model, a possible explanation of their disagreement for $w_Q > -1$ which was noted in [5].

A surprising feature of iQCDM models is that they can predict $d_L(z)$ that are uniformly larger than predicted in Λ CDM models, “simulating” $w < -1$ at the current level of the SNIa data. The energy transfer from quintessence




Figure 3: Luminosity distance–redshift ($d_L(z)$) curves for Λ CDM, a “Big Rip” equation of state $w_Q = -1.4$ (allowed by current SNIa data [5]) and iQCDM ($\alpha = 5, \beta = 15, \Omega_{Q0} = 0.6$). The inset shows the percentage difference from the Λ CDM curve. Also shown in the inset (dot-dashed) is the curve one would obtain if the quintessence and iCDM were both treated as fluids with equations of state given by their decay rate: $w = -\beta/(\alpha + \beta) = 0.75$.

to a species of matter causes that matter to have a slower decay rate. Thus one expects a *larger* luminosity distance for a given redshift than one would obtain from a model where quintessence did not interact and had a constant w_Q .

As shown in Fig. 3 for an example with ($\alpha = 5, \beta = 15, \Omega_{Q0} = 0.6$), the current equation of state of quintessence is close to -1 , and thus $d_L(z)$ is similar to a model with the same Ω_{M0} and w_Q less than -1 (though the shape is different), or nearly identical to a non-interacting model with $\Omega_{Q0} = \Omega_{Q0}(\beta + \alpha)/\beta$, and $w_Q = -\beta/(\beta + \alpha)$. As SNIa observations improve in both number and redshift depth, if we measure a larger luminosity distance than in a Λ CDM model we have the choice between two possible conclusions: ρ_Q increases with expansion as for example in theoretically problematic phantom energy models or quintessence and dark matter interact. If we can determine Ω_{M0} by other means we may be able to rule out one or the other based on the shape of $d_L(z)$. Since the physical equation of state never drops below $w = -1$, objects that are gravitationally bound today will remain

so forever.

A very important general aspect of iQCDM models is that the iCDM energy density changes differently with expansion to pure dilution. In this example, when ϕ is in the potential minimum, energy is transferred from quintessence to the iCDM. As the quintessence comes to dominate at late times, the iCDM becomes significant, and $\rho_M(\text{scale factor})^3$ increases. This could create observational signatures which have not been looked for thus far. It would thus be useful to reexamine previous experimental determinations of the matter density to place constraints on $\rho_M(\text{scale factor})^3$ at different epochs. Large-scale structure data must be reconsidered: because $\rho_M(\text{scale factor})^3$ is not constant, one must carefully consider at what epoch an observation measures it. Energy transfer and the fifth-force will have significant effects on structure formation in iQCDM effects. Detailed calculations of structure formation in these models are ongoing. Preliminary results show that there are combinations of the parameters in our model that fit observations of the cosmic microwave background as well or better than standard models [21].

To summarize: we present a class of models where we allow couplings of the dark energy with dark matter degrees of freedoms. Within these models the same condition that avoids early cosmic acceleration also makes the quintessence equation of state more negative today. Note that the parameters of the Lagrangian only need to satisfy inequalities for these results to hold, obviating the need for fine tuning. If our dark sector is iQCDM-like, we would predict a larger luminosity distance to a given redshift than for non-interacting dark energy with the same constant equation of state. Therefore current indications of $w < -1$ can be re-interpreted in our scenario, removing the impending doom of a “Big Rip” in the near cosmic future.

Acknowledgments

The authors would like to thank R. Caldwell, S. Carroll, P. Ricker, and P. Steinhardt for stimulating discussions. BDW is a 2003/4 NCSA Faculty Fellow.

-
- [1] S. M. Carroll, astro-ph/0301342
 - [2] A. G. Riess et al., AJ **116**, 1009 (1998)
 - [3] S. Perlmutter, et al., ApJ **517**, 565 (1999) 565-586
 - [4] S. Perlmutter, M. S. Turner, M. White, Phys. Rev. Lett. **83**, 670-673 (1999)
 - [5] R. A. Knop, et al., ApJ, in print astro-ph/0309368
 - [6] A. G. Riess, et al., astro-ph/0402512
 - [7] Y. Wand, M. Tegmark, astro-ph/0403292
 - [8] Wetterich, Astron. Astrophys. 301 (1995) 321-328; P. J. E. Peebles, Bharat Ratra, Rev. Mod. Phys. **75** (2003) 559-606; Ivaylo Zlatev, Limin Wang, and Paul J. Steinhardt, Phys.Rev.Lett. **82** (1999) 896-899
 - [9] E. J. Copeland, A. R. Liddle, and D. Wands, Phys. Rev. D **57** (1998) 4686-4690
 - [10] P. J. Steinhardt, L. Wang, and I. Zlatev, Phys. Rev. D **59** (1999) 123504
 - [11] A. Albrecht, C. Skordis, Phys. Rev. Lett. **84**, 2076 (2000)
 - [12] R. Caldwell Phys. Lett. **B545**, 23 (2002)
 - [13] S. M. Carroll, M. B. Hoffman, and M. Trodden, Phys. Rev. D **68**, 023509 (2003)
 - [14] R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, Phys. Rev. Lett. **91**, 071301 (2003)
 - [15] G. Farrar and J. Peebles, astro-ph/0307316
 - [16] M. B. Hoffman, astro-ph/0307350

- [17] G. W. Anderson, S. M. Carrol, astro-ph/9711288; J. A. Casas, J. Garcia-Bellido, and M. Quiros, *Class. Quantum Grav.* **9**, 1371 (1992); L. Amendola, *Phys. Rev. D* **62**, 043511 (2000); L. Amendola, D. Tocchini-Valentini, *Phys. Rev. D* **64**, 043509 (2001); L. Amendola, *Mon. Not. Roy. Astron. Soc.* **342**, 221 (2003); M. Pietroni, *Phys. Rev. D* **67**, 103523 (2003); D. Comelli, M. Pietroni, and A. Riotto, *Phys. Lett. B* **571**, 115 (2003)
- [18] Urbano Franca, Rogerio Rosenfeld, astro-ph/0308149
- [19] P. G. Ferreira, M. Joyce, *Phys. Rev. Lett.* **79**, 4740 (1997); *Phys. Rev. D* **58**, 023503 (1998).
- [20] G. Huey, R. Tavakol, *Phys. Rev. D* **66** (2002) 043514
- [21] G. Huey and B.D. Wandelt, in preparation
- [22] Briefly, one solves the equation of motion of χ as if ϕ were a constant, and then substitutes that solution for χ in the ϕ equation of motion. The result is the effective equation of motion for ϕ , and the potential appearing in the Lagrangian from which this effective equation of motion would be derived is the effective potential V .